



POST-OPERATIVE INSTRUCTIONS: DISCHARGE INSTRUCTIONS FOR JOINT REPLACEMENTS

1. Wound care

- a. Remove adhesive Silverlon dressing after 7 days
- b. If the wound is dry, you may leave it open to air or cover it with clean gauze
- c. DO NOT USE OINTMENTS OR CREAMS ON THE WOUND (SUCH AS NEOSPORIN)
- 2. DVT prophylaxis (blood clot prevention)
 - a. ASA 81 mg twice a day for 30 day (unless otherwise instructed)
 - Knee high TED hose on both legs at all times except bathing for 4 weeks

3. Ice

 a. Ice pack to operative site 4 times daily minimum 20 mins each and after PT

4. Bathing

- a. You may not shower with the Silverlon dressing in place
- b. Once the Silverlon dressing is removed and the wound is dry, you may shower. Dry the incision with a clean towel. No baths or swimming until the staples are removed.

5. Pain medications

- a. Oxycodone 5 mg 1 or 2 pills every 4 hours as needed
- b. Tylenol 1000 mg 3x/day
- c. Celebrex or Mobic as directed

6. Follow up in office

- a. 2 weeks to remove staples and wound check, appointment with PA-C
- b. 6 weeks for x-ray and to evaluate progress, appointment with Physician

7. Notify our office immediately if:



William A. Tyndall, M.D., Ph.D. 3000 Fairway Drive, Altoona, PA 16602 814-231-2101

- a. Temperature greater than 101 degrees (check with a thermometer)
- b. Worsening wound drainage, erythema or redness
- c. Readmission to hospital
- 8. Physical therapy on discharge
 - a. Home exercises per attached sheet
- 9. Dr. Tyndall can be reached at 814-574-1855
 - a. The office can be reached at 814-942-1166 anytime
 - b. If you experience an emergency go to the ER or call 911